





ZytoLight®


Products for FISH analysis

Signal Interpretation Guide

Zyto Light® SPEC ALK / EML4 TriCheck™ Probe








Normal ALK-EML4

ALK-EML4 Inversion


Non-EML4 ALK Translocation



Reliable Detection of ALK-EML4 Inversion

The **ZytoLight**® **SPEC ALK/EML4 TriCheck**™ **Probe** is designed to detect inversions [inv(2)(p21p23)] frequently observed in non-small cell lung cancer (NSCLC). These inversions are difficult to determine using conventional FISH probes, e.g. ALK dual color break apart, as the ALK and EML4 genes are mapped very close to each other. In case of an inversion, the gap between the orange and the green signal is often very small. Especially in these borderline cases, uncertainty in signal interpretation can be overcome by the ability to simultaneously check for the EML4 rearrangement status using the **SPEC ALK/EML4 TriCheck**™ **Probe**. Consequently, ZytoVision's **ZytoLight**® **SPEC ALK/EML4 TriCheck**™ **Probe** is the only FISH probe leading to a reliable discrimination of three different ALK gene conditions: non-rearranged, EML4-ALK inversion, non-EML4 ALK translocation!

Zyto*Light* ® SPEC ALK/EML4 TriCheck™ Probe Design



New ALK/EML4 TriCheck™ Evaluation Criteria

For the new evaluation criteria 374 lung adenocarcinomas were investigated.

ALK rearrangement was defined as:

- ALK break apart (one green/one orange) ≥ 1 < 2 signal diameter with orange/aqua co-localization
- ALK break apart ≥ 2 signal diameter
- Isolated orange signals


Determine ALK status according to the new and simplified scoring guidelines shown in the scheme at the right published by Schildhaus HU, et al. 2016*.

New Evaluation Procedure

- 1. Localize the invasive component of a non-small cell lung cancer specimen on a corresponding H&E or IHC slide.
- 2. The area for counting should include clearly distinguishable and well distributed nuclei.
- 3. Count just **20 consecutive and non-overlapping cells** in an area of a population of tumor cells in the invasive component of the tumor.
- 4. Determine the ALK status according to the **new ALK Signal Interpretation Guide** and the **new Scoring Guidelines**Scheme.
- 5. Report if ALK status is indeterminate due to e.g. artifacts, analytic testing failure, etc. or if ALK status is discordant with other histopathologic findings and repeat test with another specimen.

The validation of FISH probes is required for each type of tissue that is intended to be tested in clinical practice since different tissue types exhibit different cell types with different nuclei diameters which may result in different cut off values. In order to correctly interpret the results, the user must validate this product prior to use in diagnostic procedures according to national and/or international guidelines.


New Scoring Guidelines for a simplified ALK FISH Analysis published by Schildhaus HU, et al. 2016



Conclusion

- The new ZytoLight® SPEC ALK/EML4 TriCheck™ Probe scoring guidelines resulted in a sensitivity and specificity of 100% compared to the Vysis LSI ALK Dual Color Break Apart Probe.
- 99% of all lung cancer specimens needed an analysis of as little as 20 tumor cells.
- Expenditure of time per case was halved.
- For cases with low tumor cell content reliable results can be obtained.

Signal Interpretation Guide



Zyto Light ® SPEC ALK/EML4 TriCheck™ Probe

Prod. No.	Product Lab	el	Tests* (Volume)	
Z-2117-50	Zyto Light SPEC ALK/EML4 TriCheck Probe C€ IVD	/	5 (50 µl)	
Z-2117-200	Zyto Light SPEC ALK/EML4 TriCheck Probe C€ IVD	/	20 (200 µl)	
Related Products				
Z-2028-5	Zyto Light FISH-Tissue Implementation Kit CE IVD Incl. Heat Pretreatment Solution Citric, 150 ml; Pepsin Solution, 1 ml; Wash Buffer SSC, 210 ml; 25x Wash Buffer A, 50 ml; DAPI/DuraTect-Solution, 0.2 ml		5	
Z-2028-20	Zyto Light FISH-Tissue Implementation Kit C IND Incl. Heat Pretreatment Solution Citric, 500 ml; Pepsin Solution, 4 ml; Wash Buffer SSC, 560 ml; 25x Wash Buffer A, 100 ml; DAPI/DuraTect-Solution, 0.8 ml		20	
Z-2099-20	Zyto Light FISH-Cytology Implementation Kit C Incl. Cytology Pepsin Solution, 4 ml; 20x Wash Buffer TBS, 50 ml; 10x MgCl ₂ , 50 ml; 10x PBS, 50 ml; Cytology Stringency Wash Buffer SSC, 500 ml; Cytology Wash Buffer SSC, 500 ml; DAPI/DuraTect-Sol	ution, 0.8 m	20 I	

* Using 10 µl probe solution per test. CE IVD only available in certain countries. All other countries research use only! Please contact your local dealer for more information.

ZytoLight® FISH-Tissue Implementation Kit



Zyto Light ® Fluorochromes

- · Fluorochromes of the FISH probes
- Appropriate filter sets

Fluorochrome	Excitation	Emission	Equivalent to
 ZyBlue[™] 	418 nm	467 nm	DEAC
 ZyGreen[™] 	503 nm	528 nm	FITC
ZyOrange [™]	547 nm	572 nm	Rhodamine

Recommended Filter Sets

noise ratio and need to be assembled

Prod. No	Product	Detected Fluorochrome
E-4030-1	DAPI Single Bandpass Filter Set v2	DAPI
E-4026-1	ZyBlue™ Single Bandpass Filter Set v2	•
E-4012-1	ZyGreen™ Single Bandpass Filter Set v2	•
E-4013-1	ZyOrange™ Single Bandpass Filter Set v2	•
E-4016-1	ZyGreen™/ZyOrange™ Dual Bandpass Filter Set v2	•/•